

## A NOTE ON AN INVERSE THEOREM FOR A GENERALISED MODULUS OF SMOOTHNESS

MUHARREM Q. BERISHA AND FATON M. BERISHA

**ABSTRACT.** We prove the theorem converse to Jackson's theorem for a modulus of smoothness of the first order generalised by means of an asymmetric operator of generalised translation.

### INTRODUCTION

The relation between the modulus of smoothness and best approximation by trigonometric polynomials of a  $2\pi$ -periodic function is well-known. In the case of non periodic functions there is no such relation between their moduli of smoothness and best approximation by algebraic polynomials. An analogy with the  $2\pi$ -periodic case takes place if the ordinary modulus of smoothness is replaced by a generalised modulus of smoothness (see e.g. [1, 2, 5]).

In number of papers generalised moduli of smoothness are introduced by means of generalised symmetric operators of translation [3, 4, 5].

In [6], an asymmetric operator of generalised translation is introduced, by means of it a generalised modulus of smoothness of the first order is defined, and the theorem of coincidence of the class of functions defined by that modulus with the class of functions with given order of best approximation by algebraic polynomials is proved.

In the present paper we prove a theorem converse to Jackson's theorem related to that modulus of smoothness.

### 1. DEFINITIONS

By  $L_p$  we denote the set of functions  $f$  measurable on the segment  $[-1, 1]$  such that for  $1 \leq p < \infty$

$$\|f\|_p = \left( \int_{-1}^1 |f(x)|^p dx \right)^{1/p} < \infty,$$

and for  $p = \infty$

$$\|f\|_\infty = \operatorname{ess\,sup}_{-1 \leq x \leq 1} |f(x)| < \infty.$$

Denote by  $L_{p,\alpha}$  the set of functions  $f$  such that  $f(x)(1-x^2)^\alpha \in L_p$ , and put

$$\|f\|_{p,\alpha} = \|f(x)(1-x^2)^\alpha\|_p.$$

By  $E_n(f)_{p,\alpha}$  we denote best approximation of a function  $f \in L_{p,\alpha}$  by algebraic polynomials of degree not greater than  $n-1$ , in  $L_{p,\alpha}$  metrics, i.e.

$$E_n(f)_{p,\alpha} = \inf_{P_n} \|f - P_n\|_{p,\alpha},$$

where  $P_n$  are algebraic polynomials of degree not greater than  $n-1$ .

---

1991 *Mathematics Subject Classification.* Primary 41A35, Secondary 42A05.

*Key words and phrases.* Generalised modulus of smoothness, asymmetric operator of generalised translation, converse Jackson theorem, best approximations by algebraic polynomials.

For a function  $f$  we define an operator of generalised translation  $\hat{T}_t(f, x)$  by

$$\begin{aligned} \hat{T}_t(f, x) = \frac{1}{\pi(1-x^2)} \int_0^\pi & \left( 1 - \left( x \cos t - \sqrt{1-x^2} \sin t \cos \varphi \right)^2 \right. \\ & \left. - 2 \sin^2 t \sin^2 \varphi + 4(1-x^2) \sin^2 t \sin^4 \varphi \right) \\ & \times f(x \cos t - \sqrt{1-x^2} \sin t \cos \varphi) d\varphi. \end{aligned}$$

By means of that operator of generalised translation we define the generalised modulus of smoothness by

$$\hat{\omega}(f, \delta)_{p, \alpha} = \sup_{|t| \leq \delta} \left\| \hat{T}_t(f, x) - f(x) \right\|_{p, \alpha}.$$

Put  $y = \cos t$ ,  $z = \cos \varphi$  in the operator  $\hat{T}_t(f, x)$ , we denote it by  $T_y(f, x)$  and rewrite it in the form

$$\begin{aligned} T_y(f, x) = \frac{1}{\pi(1-x^2)} \int_{-1}^1 & (1 - R^2 - 2(1-y^2)(1-z^2) \\ & + 4(1-x^2)(1-y^2)(1-z^2)^2) f(R) \frac{dz}{\sqrt{1-z^2}}, \end{aligned}$$

where  $R = xy - z\sqrt{1-x^2}\sqrt{1-y^2}$ .

By  $P_\nu^{(\alpha, \beta)}(x)$  ( $\nu = 0, 1, \dots$ ) we denote the Jacobi's polynomials, i.e. algebraic polynomials of degree  $\nu$  orthogonal with the weight function  $(1-x)^\alpha(1+x)^\beta$  on the segment  $[-1, 1]$  and normed by the condition  $P_\nu^{(\alpha, \beta)}(1) = 1$  ( $\nu = 0, 1, \dots$ ).

Denote by  $a_n(f)$  the Fourier-Jacobi coefficients of a function  $f$ , integrable with the weight function  $(1-x^2)^2$  on the segment  $[-1, 1]$ , with respect to the system of Jacobi polynomials  $\{P_n^{(2,2)}(x)\}_{n=0}^\infty$ , i.e.

$$a_n(f) = \int_{-1}^1 f(x) P_n^{(2,2)}(x) (1-x^2)^2 dx \quad (n = 0, 1, \dots).$$

The following properties of the operator  $T_y$  are proved in [6].

**Lemma 1.1.** *Operator  $T_y$  has the following properties*

- (1) *The operator  $T_y(f, x)$  is linear with respect to  $f$ ;*
- (2)  *$T_1(f, x) = f(x)$ ;*
- (3)  *$T_y(P_n^{(2,2)}, x) = P_n^{(2,2)}(x) R_n(y)$  ( $n = 0, 1, \dots$ ),*  
*where  $R_n(y) = P_{n+2}^{(0,0)}(y) + \frac{3}{2}(1-y^2) P_n^{(2,2)}(y)$ ;*
- (4)  *$T_y(1, x) = 1$ ;*
- (5)  *$a_k(T_y(f, x)) = R_k(y) a_k(f)$  ( $k = 0, 1, \dots$ ).*

**Lemma 1.2.** *Let given numbers  $p, \alpha, r$  and  $\lambda$  be such that  $1 \leq p \leq \infty$ ,  $0 < \lambda < 2$ ;*

$$\begin{aligned} \frac{1}{2} < \alpha \leq 1 & \quad \text{for } p = 1, \\ 1 - \frac{1}{2p} < \alpha < \frac{3}{2} - \frac{1}{2p} & \quad \text{for } 1 < p < \infty, \\ 1 \leq \alpha < \frac{3}{2} & \quad \text{for } p = \infty. \end{aligned}$$

*Let  $f \in L_{p, \alpha}$ . Then*

$$E_n(f)_{p, \alpha} \leq \frac{C_1}{n^\lambda}$$

*if and only if*

$$\hat{\omega}(f, \delta)_{p, \alpha} \leq C_2 \delta^\lambda,$$

where constants  $C_1$  and  $C_2$  do not depend on  $f$ ,  $n$  and  $\delta$ .

The lemma is proved in [6].

## 2. THE CONVERSE THEOREM

Now we formulate our result.

**Theorem 2.1.** *Let given numbers  $p$ ,  $\alpha$  and  $\lambda$  be such that  $1 \leq p \leq \infty$ ,  $0 < \lambda < 2$ ;*

$$\begin{aligned} \frac{1}{2} < \alpha &\leq 1 && \text{for } p = 1, \\ 1 - \frac{1}{2p} < \alpha &< \frac{3}{2} - \frac{1}{2p} && \text{for } 1 < p < \infty, \\ 1 &\leq \alpha < \frac{3}{2} && \text{for } p = \infty. \end{aligned}$$

If  $f \in L_{p,\alpha}$ , then the following inequality holds

$$\hat{\omega}\left(f, \frac{1}{n}\right)_{p,\alpha} \leq \frac{C_1}{n^2} \sum_{\nu=1}^n \nu E_{\nu}(f)_{p,\alpha},$$

where the constant  $C$  does not depend on  $f$  and  $n$ .

*Proof.* Let  $P_n(x)$  be the polynomial of degree not greater than  $n - 1$  such that

$$\|f - P_n\|_{p,\alpha} = E_n(f)_{p,\alpha} \quad (n = 1, 2, \dots),$$

and

$$Q_k(x) = P_{2^k}(x) - P_{2^{k-1}}(x) \quad (k = 1, 2, \dots),$$

$$Q_0(x) = P_1(x).$$

For given  $n$  we chose the positive integer  $N$  such that

$$\frac{n}{2} < 2^N \leq n + 1.$$

By the proof of Lemma 1.2 given in [6] it follows that

$$\begin{aligned} \hat{\omega}\left(f, \frac{1}{n}\right)_{p,\alpha} &\leq C_2 \left( E_{2^N}(f)_{p,\alpha} + \frac{1}{n^2} \sum_{\mu=1}^N 2^{2\mu} \|Q_k\|_{p,\alpha} \right) \\ &\leq 2C_2 \left( E_{2^N}(f)_{p,\alpha} + \frac{1}{n^2} \sum_{\mu=1}^N 2^{2\mu} \left( E_{2^\mu}(f)_{p,\alpha} + E_{2^{\mu-1}}(f)_{p,\alpha} \right) \right) \\ &\leq 4C_2 \left( E_{2^N}(f)_{p,\alpha} + \frac{1}{n^2} \sum_{\mu=0}^{N-1} 2^{2(\mu+1)} E_{2^\mu}(f)_{p,\alpha} \right) \\ &\leq \frac{C_3}{n^2} \sum_{\mu=0}^N 2^{2(\mu+1)} E_{2^\mu}(f)_{p,\alpha}. \end{aligned}$$

Considering that for  $\mu \geq 1$  we have

$$\sum_{\nu=2^{\mu-1}}^{2^\mu-1} \nu E_{\nu}(f)_{p,\alpha} \geq E_{2^\mu}(f)_{p,\alpha} 2^{2(\mu-1)},$$

it follows that

$$\begin{aligned}\hat{\omega}\left(f, \frac{1}{n}\right)_{p,\alpha} &\leq \frac{C_4}{n^2} \left(2^2 E_1(f)_{p,\alpha} + \sum_{\mu=1}^N \sum_{\nu=2^{\mu-1}}^{2^{\mu}-1} \nu E_{\nu}(f)_{p,\alpha}\right) \\ &\leq \frac{C_5}{n^2} \sum_{\nu=1}^n \nu E_{\nu}(f)_{p,\alpha}.\end{aligned}$$

Theorem 2.1 is proved.  $\square$

#### REFERENCES

1. P. L. Butzer, R. L. Stens, and M. Wehrens, *Approximation by algebraic convolution integrals*, Approximation theory and functional analysis (Proc. Internat. Sympos. Approximation Theory, Univ. Estadual de Campinas, Campinas, 1977), North-Holland, Amsterdam, 1979, pp. 71–120. MR 81h:41021
2. Z. Ditzian and V. Totik, *Moduli of smoothness*, Springer-Verlag, New York, 1987. MR 89h:41002
3. M. K. Potapov, *O strukturnykh kharakteristikakh klassov funktsii s dannym poryadkom nailuchshego priblizheniya*, Trudy Mat. Inst. Steklov. **134** (1975), 260–277, 410. MR 53 #6184
4. ———, *Ob usloviyakh sovpadeniya nekotorykh klassov funktsii*, Trudy Sem. Petrovsk. (1981), no. 6, 223–238. MR 82i:46053
5. ———, *O priblizhenii algebraicheskimi mnogochlenami v integral'noi metrike s vesom Yakobi*, Vestnik Moskov. Univ. Ser. I Mat. Mekh. (1983), no. 4, 43–52. MR 84i:41008
6. ———, *O sovpadenii klassov funktsii opredelyaemykh operatorom obobshchennogo sdviga ili poryadkom nailuchshego priblizheniya algebraicheskimi mnogochlenami*, Mat. Zametki **66** (1999), no. 2, 242–257. MR 2000k:41008

F. M. BERISHA, FACULTY OF MATHEMATICS AND SCIENCES, UNIVERSITY OF PRISHTINA, NËNA TEREZË 5, 10000 PRISHTINË, KOSOVË  
E-mail address: faton.berisha@uni-pr.edu